

OVARIAN CANCER

SYMPTOMS

- ▶ Bloating
- ▶ Pelvic or abdominal pain
- ▶ Difficulty eating or feeling full quickly
- ▶ Urinary symptoms (urgency or frequency)



SURGERY
IS REQUIRED
TO ACCURATELY STAGE
OVARIAN CANCER.

DEVELOPMENT PIPELINE

Learn more in ISR's
"Epithelial Ovarian
Cancer: Disease
and Pipeline
Analysis"
on ISRreports.com

TREATMENT



surgery



chemotherapy



targeted
therapy

RISK FACTORS



GENETICS:
BRCA1 and BRCA2
genes, Lynch
Syndrome, and
family history

**INCREASING AGE
REPRODUCTIVE HISTORY
& INFERTILITY**



**HORMONE
REPLACEMENT THERAPY**

1 IN 77


WOMEN WILL BE DIAGNOSED
WITH OVARIAN CANCER
DURING THEIR LIFETIMES.

PROGNOSIS


44.6%

AVG. 5 YEAR SURVIVAL RATE

PERCENT OF CASES BY STAGE



5 YEAR SURVIVAL



PREVALENCE:

In 2011, there were approximately
188,867 WOMEN

living with ovary cancer in the
United States.

That's about the population of
Tallahassee, FL.

1.3%

of women are
diagnosed with
ovarian cancer
in their
lifetimes



**INDIA, THE
UNITED STATES,
AND CHINA**

have the highest number
of annual deaths from
Ovarian cancer



63

Median
Age At
Diagnosis